

Buffer contribution to formation enthalpy of copper(II)–bicine complex determined by isothermal titration calorimetry method

A. Tesmar¹ · D. Wyrzykowski¹ · D. Jacewicz¹ · K. Żamojć¹ · J. Pranczk¹ ·
L. Chmurzyński¹

Received: 14 October 2015 / Accepted: 26 January 2016 / Published online: 23 February 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract The isothermal titration calorimetry (ITC) technique supported by potentiometric titration data was used to study a buffer contribution to the formation enthalpy of copper(II)–bicine complex obtained directly from the ITC experiments. The calorimetric measurements were carried out at 298.15 K in 100 mM buffer solutions with a pH of 6. In experiments two biologically relevant pH buffers were used, namely Mes and Caco. These buffers do not bind copper(II) ions and thus do not affect the binding constant. However, due to their different ionization enthalpy, they affect the conditional (observed) enthalpy of binding of the metal ion to a ligand owing to the proton transfer during the interactions. To calculate the conditional-independent binding enthalpy ΔH , the equation based on the Hess's law was used. Furthermore, pH-independent and buffer-independent parameters (K , ΔG and ΔS) of the interactions were calculated as well. The relationship between the ionization enthalpy of the buffer components and the thermodynamic parameters has been discussed.

Keywords Cu(II)–bicine complex · Buffer · Proton transfer · Thermodynamic parameters · Isothermal titration calorimetry

Introduction

Isothermal titration calorimetry (ITC) measurements are usually carried out in buffer solutions. Such an approach enables to eliminate a pH mismatch between a titrant (a syringe solution) and a titrand (a cell solution) and thus prevents a generation of an additional heat on account of a neutralization reaction ($\text{H}_3\text{O}^+ + \text{OH}^- = 2\text{H}_2\text{O}$) that is not connected with molecular interactions [1–4]. Furthermore, the proper maintenance of the pH is of a particular importance when studying systems of biological interest, for instance proteins (the change of pH modifies electrostatic interactions between charge functional groups of the amino acids and consequently the three-dimensional structure of the protein) [5] or processes involving enzymatic reactions that can only take place in a narrow range of pH [6]. It is also worth noticing that the choice of an appropriate buffer is of great importance as it has an impact on binding constants as well as the enthalpy of complex formation, especially for those complexes containing metal ions [5, 7–9]. If the buffer with an affinity for a metal ion is used, it affects the metal–ligand binding constant. Therefore, to obtain buffer-independent parameters (K and ΔG) the metal–buffer stability constants need to be available (or determined) and subsequently taken into account during the data analysis [10, 11]. Moreover, if a process of complex formation is accompanied with proton transfers, the equivalent number of protons is taken up or released by the buffer. It generates an additional heat that is proportional to the enthalpy of buffer ionization [12, 13]. In this case, the enthalpy measured during the ITC experiment reflects both the buffer ionization and complex formation [14].

In this article, we present the influence of the type of a buffer (the ionization enthalpy of the buffer) on the

✉ D. Wyrzykowski
dariusz.wyrzykowski@ug.edu.pl

¹ Faculty of Chemistry, University of Gdańsk, Wita Stwosza 63, 80-308 Gdańsk, Poland

thermodynamic parameters for the binding of Cu^{2+} to bicine [*N,N*-bis(2-hydroxyethyl)glycinate] that is used in these studies as a ligand (Fig. 1). In experiments 2-(*N*-morpholino)ethanesulfonic acid (Mes) and dimethylarsenic acid (Caco) were used as buffer substances that differ in their ionization enthalpies ($\text{BH}^{\pm} = \text{B}^{-} + \text{H}^{+}$), +3.54 and $-0.72 \text{ kcal mol}^{-1}$ for Mes and Caco, respectively [15].

It is also worth emphasizing that bicine is also a useful buffer standard in the range of physiological interest [16, 17] with the ionization enthalpy equal to $-6.47 \text{ kcal mol}^{-1}$ [15]. It has also been employed in colour photographic processes, in analytical methods, as a stabilizing agent. Bicine acts as a strong binding ligand forming a fairly stable complex with most metal ions [18]. Based on its complexation properties, the use of bicine buffer is not advisable in environmental and biological studies involving metals, unless thermodynamic parameters of the metal–bicine interaction are taken into consideration.

Experimental

Materials

All reagents, namely 2-(*N*-morpholino)ethanesulfonic acid hydrate ($\geq 99\%$) (Mes), dimethylarsenic acid ($\geq 99\%$) (Caco), $\text{Cu}(\text{NO}_3)_2 \cdot 2.5\text{H}_2\text{O}$ ($\geq 99.99\%$) and *N,N*-bis(2-hydroxyethyl)glycine ($\geq 99\%$) [$\text{H}(\text{bicine})^{\pm}$] were purchased from Sigma-Aldrich Chemical Corp.

Isothermal titration calorimetry (ITC)

All ITC experiments were performed at 298.15 K using an AutoITC isothermal titration calorimeter (MicroCal Inc., Malvern, Northampton, USA) with a 1.4491-mL sample and the reference cells. The reference cell contained distilled water. The data, specifically the heat normalized per mole of injectant, were processed with Origin 7 from MicroCal. An initial $2 \mu\text{L}$ injection was discarded from each data set in order to remove the effect of titrant diffusion across the syringe tip during the equilibration process. All reagents were dissolved directly in the 100 mM

buffer solution of Mes or Caco. The pH of the buffer solution was adjusted to 6.0 with 0.1 M HNO_3 . The experiment consisted of injecting $10.02 \mu\text{L}$ (29 injections, $2 \mu\text{L}$ for the first injection only) of ca 1.2 mM buffered solution of *N,N*-bis(2-hydroxyethyl)glycine, $\text{H}(\text{bicine})$, into the reaction cell which initially contained 0.07 mM buffered solution of $\text{Cu}(\text{NO}_3)_2$. A background titration, consisting of an identical titrant solution but with the buffer solution in the reaction cell only, was removed from each experimental titration to account for the heat of dilution. All the solutions were degassed prior to the titration. The titrant was injected at 5-min intervals to ensure that the titration peak returned to the baseline before the next injection. Each injection lasted 20 s. For homogeneous mixing in the cell, the stirrer speed was kept constant at 300 rpm. A calibration of the AutoITC calorimeter was carried out electrically by using electrically generated heat pulses. The CaCl_2 –EDTA titration was performed to check the apparatus, and the results (n —stoichiometry, K , ΔH) were compared with those obtained for the same samples (test kit) at MicroCal.

Potentiometric measurements

Potentiometric titrations were performed in the 30 mL thermostated ($298.15 \pm 0.10 \text{ K}$) cell using the Cerko Lab System microtitration unit fitted with the 0.5-mL Hamilton's syringe, pH combined electrode (Schott–BlueLine 16 pH type) and a self-made measuring cell equipped with a magnetic stirrer. The temperature was controlled using the Lauda E100 circulation thermostat. The electrode was calibrated according to IUPAC recommendations [19]. The syringe was calibrated by a weight method. All the solutions were prepared immediately before measurements. The compositions of the titrand solutions used in the experiments were as follows: (1) $\text{H}(\text{bicine})$ (2.05 mM) and HNO_3 (2.48 mM) and (2) Cu^{2+} (1.50 mM), $\text{H}(\text{bicine})^{\pm}$ (2.05 mM) and HNO_3 (2.48 mM). The solutions were potentiometrically titrated with a standardized KOH solution (25.02 mM) with the pH ranging from 2.5 to 12.0. The stability constants of the complexes were determined using the CVEQUID program [20] based on a minimization of the differences between the

Fig. 1 Proposed binding mode of copper(II) ion to *N,N*-bis(2-hydroxyethyl)glycinate (L)






Table 1 Individual reactions that contribute to the overall equilibrium for the formation of the copper(II)–bicine complex in the Mes and Caco buffer solution (B)

No.	Reaction	$\Delta H/\text{kcal mol}^{-1}$ (for the association reactions)
1	$\text{H}(\text{bicine})^{\pm} = (\text{bicine})^{-} + \text{H}^{+}$	$-\Delta_{\text{H}(\text{bicine})}H^{\circ} = -6.47$
2	$\text{Cu}^{2+} + (\text{bicine})^{-} = \text{Cu}(\text{bicine})^{+}$	$\Delta_{\text{Cu}(\text{bicine})}H^{\circ} = ?$
3	$\text{B}^{-} + \text{H}^{+} = \text{HB}^{\pm}$	$\Delta_{\text{BH}}H^{\circ} = -3.54$ (Mes) and $+0.72$ (Caco)

**Fig. 2** Plot of the pH versus the $\text{NaOH}/\text{H}^{+} [\text{HNO}_3 + \text{H}(\text{bicine})]$ molar ratio during the potentiometric titration of the $\text{H}(\text{bicine})\text{--HNO}_3$ (square) and $\text{Cu}(\text{II})\text{--H}(\text{bicine})\text{--HNO}_3$ (circle) solutions with the NaOH solution**Fig. 3** Concentration distribution curves of the complexes as a function of pH calculated based on the stability constants obtained from PT measurements

theoretical model and the experimental data, according to the Gauss–Newton–Marquart method for nonlinear equations (see ref [21] for more details).

Results and discussion

Bicine [*N,N*-bis(2-hydroxyethyl)glycinate]] acts as a tetradentate ligand and forms 1:1 metal–ligand complexes. In a solid state, one carboxylate oxygen atom, two hydroxyl oxygen atoms and one nitrogen atom participate in the copper(II) binding. In solutions an aqua ligand coordinates to the metal centre, hence completing the coordination sphere (Fig. 1).

The stability of the resulting complex (CuL , where L denotes bicine) in a solution is reflected by a value of stability constant. As far as the ITC technique is concerned, this parameter is quantified by Eq (1):

$$K_{\text{ML}} = K_{\text{ITC}} \left(1 + \frac{[\text{H}^{+}]}{K_{\text{a2}}} + \frac{[\text{H}^{+}]^2}{K_{\text{a2}} \cdot K_{\text{a1}}} \right) (1 + K_{\text{MB}}[\text{B}]) \quad (1)$$

where K_{ML} is the pH-independent and buffer-independent metal–ligand binding constant, K_{ITC} is the Cu–L conditional stability constant obtained directly from the ITC experiment, K_{a} is the proton dissociation constant of a ligand (here K_{a1} and K_{a2} of bicine), K_{MB} is the stability constant for the metal–buffer complex, and $[\text{B}]$ is the concentration of the buffer solution.

To eliminate the impact of buffer components on the thermodynamic parameters, the Mes and Caco buffers have been chosen. These buffers do not reveal an affinity towards the copper(II) ions [22–24]; therefore, they do not affect the copper(II)–ligand binding constant. For this reason, K_{ITC} is not conditioned by the buffer competition with the ligand (bicine) for the metal (Cu^{2+}) but it depends only on the proton competition with the metal for the ligand (Eq. 2):

$$K_{\text{ML}} = K_{\text{ITC}} \left(1 + \frac{[\text{H}^{+}]}{K_{\text{a2}}} + \frac{[\text{H}^{+}]^2}{K_{\text{a2}} \cdot K_{\text{a1}}} \right) \quad (2)$$

The individual equilibria that contribute to the overall equilibrium also affect the enthalpy measured directly in the ITC experiments. In Table 1 the individual equilibria taking place in the complex formation are collected. Equilibria are written in the direction that the reaction occurs.




Fig. 4 Calorimetric titration isotherms of the binding interaction between Cu^{2+} and bicine in 100 mM Mes and Caco buffers of pH 6, at 298.15 K

Table 2 Thermodynamic parameters of copper(II) binding to bicine in buffer solutions (100 mM Mes and Caco) of pH 6, at 298.15 K

Parameter	Buffer	
	Mes	Caco
$\log K_{\text{ITC}}/\text{M}^{-1\text{a}}$	5.60 (± 0.07)	5.40 (± 0.05)
$\log K/\text{M}^{-1\text{b}}$	9.02 (± 0.07)	8.82 (± 0.05)
$\Delta_{\text{ITC}}H/\text{kcal mol}^{-1\text{a}}$	-3.48 (± 0.04)	-1.86 (± 0.04)
$\Delta H/\text{kcal mol}^{-1\text{c}}$	-4.56 (± 0.04)	-4.57 (± 0.04)
$T\Delta_{\text{ITC}}S/\text{kcal mol}^{-1\text{d}}$	4.16 (± 0.07)	5.51 (± 0.07)
$T\Delta S/\text{kcal mol}^{-1\text{d}}$	7.75 (± 0.10)	7.46 (± 0.10)
$\Delta_{\text{ITC}}G/\text{kcal mol}^{-1\text{d}}$	-7.64 (± 0.10)	-7.37 (± 0.07)
$\Delta G/\text{kcal mol}^{-1\text{d}}$	-12.31 (± 0.10)	-12.03 (± 0.07)

^a The binding constant K_{ITC} and binding enthalpy $\Delta_{\text{ITC}}H$ were obtained from ITC experiments by fitting binding isotherms, using nonlinear least-squares procedures, to a model that assumes a single set of identical binding sites

^b The logarithms of Cu(II)-bicine binding constants corrected for a proton competition with the metal ion for the ligand (Eq. 2)

^c The enthalpies of Cu(II)-bicine interactions based on Eq. 3

^d $\Delta_{\text{ITC}}G$, ΔG , $T\Delta_{\text{ITC}}S$ and $T\Delta S$ calculated using the standard thermodynamic relationships: $\Delta G = -RT\ln K = \Delta H - T\Delta S$

Taking into consideration the fact that the heat absorbed or released during ITC experiments is equal to the sum of all the energetic effects corresponding to the particular

equilibria taking place in the complexation process (Table 1), the pH- and buffer-independent enthalpy of the complex formation can be calculated from the equation based on the Hess's law (Eq. 3) [25–27]:

$$\Delta_{\text{ITC}}H = n_{\text{H}^+}\Delta_{\text{BH}}H^\circ + (\Delta_{\text{Cu}(\text{bicine})}H^\circ - n_{\text{H}^+}\Delta_{\text{H}(\text{bicine})}H^\circ) \quad (3)$$

where n_{H^+} is the number of protons transferred during the complex formation.

The number of protons transferred was determined experimentally from potentiometric titration (PT) measurements according to the procedure described in the literature [9, 28]. Thus, the solutions containing $\text{H}(\text{bicine})^\pm$ and HNO_3 in the molar ratio of 1:1.21 (solution 1) and another containing Cu(II), $\text{H}(\text{bicine})$ and HNO_3 in the molar ratio of 1:1.37:1.65 (solution 2) were titrated with a standardized NaOH solution. A relationship between the pH and the $\text{NaOH}/\text{H}^+ [\text{HNO}_3 + \text{H}(\text{bicine})^\pm]$ molar ratio is presented in Fig. 2. The difference between the number of moles of NaOH, expended for neutralization of solutions 1 and 2 up to a pH of 6, corresponds to the number of protons (n_{H^+}) lost by $\text{H}(\text{bicine})^\pm$ upon a complexation of the copper(II) ions. The number of protons determined in this way is 0.39, and this value was used for calculating $\Delta_{\text{Cu}(\text{bicine})}H^\circ$ based on Eq. 3.

Furthermore, the following equilibria were used to calculate the copper(II)–bicine stability constant from PT measurements:

1. $\text{HL}^{\pm} = \text{H}^{+} + \text{L}^{-} \quad pK_{a2}$
2. $\text{Cu}^{2+} + \text{L}^{-} = \text{CuL}^{+} \quad \log K$
3. $\text{CuL}^{+} + \text{H}_2\text{O} = \text{CuL}(\text{OH}) + \text{H}^{+} \quad pK_{\text{CuL}(\text{OH})}$

The above equilibrium model has given the best fitting of the calculated data to the experimental ones. The parameters obtained based on the assumed equilibrium model are as follows (the standard deviation in parentheses): $pK_{a2} = 9.45$ (± 0.06), $\log K = 9.00$ (± 0.09) and $pK_{\text{CuL}(\text{OH})} = 6.52$ (± 0.05). Species distributions as a function of pH are shown in Fig. 3. The complexation of the Cu(II) ions starts with the formation of the binary complex $[\text{Cu}(\text{bicine})]^{+}$. The complex predominates in the solution in the wide pH range (Fig. 3). Above pH 11 the stability of the complex is not favoured over the resulting hydroxo complex, $\text{CuL}(\text{OH})$.

Representative binding isotherms for Cu^{2+} –bicine interactions in the buffers (Mes, Caco) are shown in Fig. 4, whereas conditional parameters (marked by the subscript ITC) as well as pH-independent and buffer-independent parameters of the interactions are summarized in Table 2.

Alternatively, the $n_{\text{H}^{+}}$ value can be calculated from the slope of the relationship described by Eq. 3 ($y = ax + b$). The $n_{\text{H}^{+}}$ value determined in this way ($n_{\text{H}^{+}} = 0.39$) is in good agreement with that obtained directly from PT measurements. The numerical value of the binding enthalpy calculated based on Eq. 3 is negative (Table 2). This indicates that the endothermic effects connected with the dehydration of the $[\text{Cu}(\text{H}_2\text{O})_5]^{2+}$ ion are overcompensated by the exothermic effect due to the formation of new Cu^{2+} –bicine bonds (Fig. 1).

Conclusions

Isothermal titration calorimetry (ITC) experiments supported by potentiometric titration data have successfully been applied to determine thermodynamic parameters for the complexation of the copper(II) ions with bicine. The study confirmed the fact that both Mes and Caco buffers do not bind copper(II). Thus, these buffers would be good choices for metal speciation studies within their operational pH range. It has also been presented how to include the enthalpy of proton dissociation from the ligand and the enthalpy of buffer ionization during calorimetric data analysis. These processes generate additional heat that is not connected with intermolecular interactions and should be taken into account while interpreting calorimetric data.

When a reaction involves the release (or uptake) of protons, the equation based on the Hess's law can be applied to calculate the conditional-independent binding enthalpy ΔH . Such an approach can be used provided that the number of protons ($n_{\text{H}^{+}}$) transferred during the interaction is known. We have proved that for low molecular mass ligands this value ($n_{\text{H}^{+}}$) can be easily determined using PT measurements.

Acknowledgements This project was financially supported by the National Science Centre on the basis of decision number DEC-2012/07/B/ST5/00753.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Ladbury JE, Doyle ML. Biocalorimetry 2. Application of calorimetry in the biological sciences. West Sussex, England: Wiley; 2004.
2. Gaisford S, O'Neill MAA. Pharmaceutical isothermal calorimetry. New York: Informa Healthcare USA, Inc.; 2007.
3. Salim NN, Feig AL. Isothermal titration calorimetry of RNA. Methods. 2009;47:198–205.
4. Feig AL. Applications of isothermal titration calorimetry in RNA biochemistry and biophysics. Biopolymers. 2007;87:293–301.
5. Ferreira CMH, Pinto ISS, Soares EV, Soares HMVM. (Un)suitability of the use of pH buffers in biological, biochemical and environmental studies and their interaction with metal ions—a review. RSC Adv. 2015;5:30989–1003.
6. Nielsen JE. Analyzing enzymatic pH activity profiles and protein titration curves using structure-based pKa calculations and titration curve fitting. Methods Enzymol. 2009;454:233–58.
7. Zhang Y, Akilesh S, Wilcox DE. Isothermal titration calorimetry measurements of Ni(II) and Cu(II) binding to His, GlyGlyHis, HisGlyHis, and bovine serum albumin: a critical evaluation. Inorg Chem. 2000;39:3057–64.
8. Grosseohme NE, Akilesh S, Guerinot ML, Wilcox DE. Metal-binding thermodynamics of the histidine-rich sequence from the metal-transport protein IRT1 of Arabidopsis thaliana. Inorg Chem. 2006;45:8500–8.
9. Wyrzykowski D, Zarzeczkańska D, Jacewicz D, Chmurzyński L. Investigation of copper(II) complexation by glycylglycine using isothermal titration calorimetry. J Therm Anal Calorim. 2011;105:1043–7.
10. Wyrzykowski D, Pilarski B, Jacewicz D, Chmurzyński L. Investigation of metal–buffer interactions using isothermal titration calorimetry. J Therm Anal Calorim. 2013;111:1829–36.
11. Wyrzykowski D, Tesmar A, Jacewicz D, Pranczk J, Chmurzyński L. Zinc(II) complexation by some biologically relevant pH buffers. J Mol Recognit. 2014;27:722–6.
12. Dębowski D, Wyrzykowski D, Lubos M, Rolka K. Interactions between trypsin and its peptidic inhibitors studied by isothermal titration calorimetry (ITC). J Therm Anal Calorim. 2016;123:807–12.
13. Wyrzykowski D, Czupryniak J, Ossowski T, Chmurzyński L. Thermodynamic interactions of the alkaline earth metal ions with citric acid. J Therm Anal Calorim. 2010;102:149–54.

14. Wyrzykowski D, Chmurzyński L. Thermodynamics of citrate complexation with Mn^{2+} , Co^{2+} , Ni^{2+} and Zn^{2+} ions. *J Therm Anal Calorim*. 2010;102:61–4.
15. Goldberg RN, Kishore N, Lennen RM. Thermodynamic quantities for the ionization reactions of buffers. *J Phys Chem Ref Data*. 2002;31:231–70.
16. Good NE, Winget GD, Winter W, Connolly TN, Izawa S, Singh RMM. Hydrogen ion buffers for biological research. *Biochemistry*. 1966;4:467–77.
17. Nakon R, Krishnamoorthy CR. Free-metal ion depletion by “Good’s” buffers. *Science*. 1983;221:749–50.
18. Taha M, Khalil MM, Mohamed SA. Metal ion-buffer interactions. Complex formation of N, N-bis(2-Hydroxyethyl)glycine (Bicine) with various biologically relevant ligands. *J Chem Eng Data*. 2005;50:881–7.
19. Brandariz I, Barriada J, Vilarino T, de Vicente MS. Comparison of several calibration procedures for glass electrodes in proton concentration. *Monatsh Chem*. 2004;135:1475–88.
20. Kostrowicki J, Liwo A. A general method for the determination of the stoichiometry of unknown species in multicomponent systems from physicochemical measurements. *Comput Chem*. 1987;11:195–210.
21. Kostrowicki J, Liwo A. Determination of equilibrium parameters by minimization of an extended sum of squares. *Talanta*. 1990;37:645–50.
22. Mash HE, Chin YP. Complexation of copper by Zwitterionic Aminosulfonic (Good) buffers. *Anal Chem*. 2003;75:671–7.
23. Cereghetti GM, Negro A, Vinck E, Massimino ML, Sorgato MC, Doorslaer SV. Copper(II) binding to the human Doppel protein may mark its functional diversity from the prion protein. *J Biol Chem*. 2004;279:36497–503.
24. Yu Q, Kandegedara A, Xu Y, Rorabacher DB. Avoiding interferences from Good’s buffers: a contiguous series of noncomplexing tertiary amine buffers covering the entire range of pH 3–11. *Anal Biochem*. 1997;253:50–6.
25. Baker BM, Murphy KP. Evaluation of linked protonation effects in protein binding reactions using isothermal titration calorimetry. *Biophys J*. 1996;71:2049–55.
26. Fukada H, Takahashi K. Enthalpy and heat capacity changes for the proton dissociation of various buffer components in 0.1 M potassium chloride. *Proteins*. 1998;33:159–66.
27. Haq I, O’Brien R, Lagunavicius A, Siksnyš V, Ladbury JE. Specific DNA recognition by the type II restriction endonuclease *MunI*: the effect of pH. *Biochemistry*. 2001;40:14960–7.
28. Christensen T, Gooden DM, Kung JE, Toone EJ. Additivity and the Physical Basis of multivalency effects: a thermodynamic investigation of the calcium EDTA interaction. *J Am Chem Soc*. 2003;125:7357–66.